RESTRICTED USE PESTICIDE

Do to acute oral and inhalation toxicity. For retail sale to and application only by certified applicators or personnel under their direct supervision.

Furadan, 4F Insecticide - Nematicide

EPA SLN NO. CO-920001

FOR DISTRIBUTION AND USE ONLY IN **COLORADO**

EMERGENCY CALLS: 800-331-3148

For Agricultural or Commercial Use Only EPA Reg. No. 279-2876 ZA

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING		
CROP	PEST	RATE OF APPLICATION
Potatoes	Nematodes (aids in control)	7.2 fluid ounces per 1,000 feet of row (3 quarts per acre based on 40" rows)

DIRECTIONS FOR USE:

Use Furadan 4F at a rate of 7.2 fluid ounces per 1,000 feet of row (3 quarts per acre on 40 inch row spacing, 3.46 quarts on 34 inch spacing). Time application from planting time in-furrow to early emergence of potato plants.

Planting Time

Apply across the width of the furrow.

Preemergent Applications:

For Sprinkler Irrigated Fields:

Apply as a 4 to 6 inch band over the top of the hill in sufficient carrier for thorough coverage. Irrigate as soon as possible after treatment, but do not over irrigate. Refer to the Federal label for potential groundwater precautions.

For Furrow (Gravity) Irrigated Fields:

Shank in 4 to 6 inches to the water side of the row and approximately 4 inches deep. Irrigate as soon as possible after treatment, but do not over irrigate. Refer to the Federal label for potential groundwater precautions.

Furadan 4F may be mixed with liquid fertilizers or water. If liquid fertilizer is used, check physical compatibility before mixing large quantities.

RESTRICTIONS:

Do not make a foliar application of Furadan 4F on potatoes if Furadan has been applied at planting.

Do not exceed a rate of 7.2 fluid ounces per 1.000 feet of row per acre per season when applied under this SLN 24(c).

Do not apply more than 2 pints per acre per season in areas having greater than 90% sand and less than 1% organic matter where the water table (groundwater) is shallow (less than thirty feet).

Do not apply within 75 days of harvest.

Do not use in San Luis Valley!

For waterfowl protection, do not apply on field in proximity of waterfowl nesting areas and/or on fields where waterfowl are known to repeatedly feed.

Do not use this product through any type of irrigation system.

FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARD REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED LABEL. THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION. THIS LABEL FOR FURADAN 4F EXPIRES AND MUST NOT BE DISTRIBUTED OR USED. IN ACCORDANCE WITH THIS SLN REGISTRATION AFTER DECEMBER 31, 2007.

